AD666807

| ı |
|---|
|   |
| ı |
|   |
|   |
|   |
| 1 |
|   |

REPORT NUMBER 2

ARBOVIRUS STUDIES IN SÃO FAULO, BRASIL

ALUAL REPORT

BY

Oscar de Souna Lopes

JALUARY 1967


U.S. Army Element Defense Research Office for Latin America Rie de Jameiro, Brasil DA-HC19-67-C-C010

U.S. Department of the Army Project Number 3A014501B71P

Instituto Adolfo Lutz Laboratório de Arbovins São Paulo, Brasil

Distribution of Tric Document is Unlimited

# JISCLAIMER NOTICE



THIS DOCUMENT IS BEST QUALITY AVAILABLE. THE COPY FURNISHED TO DTIC CONTAINED A SIGNIFICANT NUMBER OF PAGES WHICH DO NOT REPRODUCE LEGIBLY.

### Introduction

The studies on the epidemiology of the Arboviruses in São Paulo, Brasil, were continued as outlined in the last year report.

The routine to isolate viruses and to collect the samples has already been described in previous reports.

### Yammes.

The agents isolated and identified this year are shown in Table I. It can be seen that 14 of them were isolated from birds captured in the study areas. The animals were netted during the period from September 66 to April 67 and by CF tests they are all identical, seeming to be different isolations of the same virus. Neutralization tests in TC are in progress.

The prototype choosen was Ar. 5245, the first isolate in the Laboratory. It was shown to be an agent sensitive more than 3 logs to DCA, with an AST of 2,7 for suchling also inoculated intracerebrally (IC) and 4.0 for those inoculated intraperitorially (IP). The virus killed abult hime inoculated IC in 9 days and those inoculated IP survived and slowed a good CF titer in their sero. The agent was filtered through Millipore Filter with membrane pore size of 450 mm.

Antigens prepared with sucrose-scetche extraction of the baby mice trains showed a herogelutin (HA) with a titer of 160 for goose cells, and a good CF antigen. Antigens of baby mice sern and liver did not show a HA antigen. This agent produce a good CFE in BHK-21 and VERO Cells.

The virus was tested in HI and CF tests against the following viruses: EEE, WEE, VEE, Mucanto, Pixuna, Mayaro, Yellow Fever (Asibi), SLE, Bussuquara, Ilheus, Caraparu, Orliver, Guaroa, Cache Valley, Maguari, California BFS, Iccaracy, Itaporanga, Anhanga, Boraceia, Juniu, Tacario, Tacaiuma, Guama, Catu, Capim, Buslara, Bush-bush, Manzan'lba, Melao, Anophelos A, Timbo, Turkeck, Coeal, Irituia, Acara, Mirin, Cotia. No cross reactions were shown. It seemed to us that we have an agent DCA sensitive, different of all existing in the Laboratory and problably an arbovirus by its behaviour. The agent was sent to Yale Arbovirus Research Enit for further studies.

Another virus, Ar 5881, was isolated from a pool of 100 Aedes servatus captured in Itapetinings on 10/11/66. It killed halv wice IC with an AST of 3.1 and adults IC with an AST of 7.0. It did not kill adults incoulated IP but a good immune sera was obtained. Antipens prepared by sucrese-acetone entraction of taby wice trains, livers and sera did not yeld an HA. In OF tests a reaction was obtained with members of California Complex, being the strongest with Melac virus. Neutralization tests are in progress to see if the virus is a isolation of Melac virus in our area of a new virus in the California Complex.

Another viruses were identified by now. Ar 2984 and Ar 2546 are the same isolation of a virus belonging to the Bunyan were Group. This

agent was different from all viruses existing in the laboratory and was sent to the WEO World Reference Center at Yale University for final ide tification. A study of the local population is being made, as some for the human sera here are able to neutralize the agent.

An 3088, Ar 2573, Ar 2494 are new strains of Cotia virus (Am.J. Trop.Med. & Hyg., 14: 156, 1965). These isolations were the first obtained in Gasa Grande of a virus isolated in another field statior. Its presence in the human population is under study.

Ar 5507 is an agent isolated from 100 A.cruzii collected on 11/5/67 in Casa Grande and it is a new isolation of Boraceia Virus, Ar 395.

Other viruses are being studied. Ar 4253, isolated from Cotia on 1/2/66 of a pool of <u>Psorophora ferox</u>, Ar 6171 from a pool of <u>Wyeo-myia confusa</u> collected in 7 cs Grande on 12/2/66 and Ar 6629 from a pool of <u>Culex</u> collected in Casa Grande on 2/3/67. They seemed to be different from the agents isolated by this Laboratory and their identification are in progress.

Ar 6629 was a very interesting isolation because was obtained from mosquitoes collected inside of the houses of the village.

### Tentative of Viral Isolation From Human Beings

In spite of the work done by us with the help of the local school teacher, we continue to face a strong resistence from the local population to accuse any illness.

We obtained samples from 2 febrile cases from children attending the school. The viral isolations were negative and the serclogy, using all the artigens that we have in the Laboratory also did not shown any rise in titer for Arbovirus.

From a small city near the border of the State of Parana we received blood from 7 children with fever, headache and signs of envolviment of nervous system. All of them were recently vaccinated against Yellow Fever. The tentative of viral isolations were negative and no rise in titer for arbovirus was observed in the serology. The serological survey made with sera collected in 1966 in Casa Grande was almost completed. The total results are shown in Table II.

It can be seen that the results are almost the same obtained in previous bleeding. Tacaiuma and Boraceia viruses are the more common in the area with a prevalence of 10% for Boraceia virus with 7 convertions to positive and 16% for Tacaiuma virus, with 14 convertions. Other agents isolated localy were also present in the population.

Ar 4175, an agent related to Boraceic virus was positive for only 9 people showing a smaller prevalence than Boraceia virus.

Ar 2984, a merber of the Bunyamwera Group was also positive in 7 sera, with prevalence smaller than Boraceia and Tacaiuma viruses.

The antibodies for the B Group were present again, in the levels obtained previously. Bussuquara was the more common virus found, but the titers were low and the pattern seemed to be a cross reaction, as observed previously. We detected 12 convertions.

### 

The following the following state of the y and y are the following y

## <u>Vertearning</u>

. The term of the of organism pass of blome in Twices IV and  $\mathbf{v}_{\bullet}$ 

The galaxy points of the period of the results are the presented factors

For struc, is then denoted by a large firing a new state of the functional form and the few Masses of the (Survey in Asia), in Figure 1 tests, As Sunt form on tests of the first sector, has been a test of the function.

### Commer.

Due to their persons Open Geneils work of decreases with the persons for the fact of the strain of Levy modern loss for article strain of the strain.

The period about one in the bleeds, a mode or Isr5 and Isr6 subject that they been periodically inflated.

In that is a less that the human holigs were indected regularly name of the holese at algority by mosquither that involve their hores. To copy of the hypothesis shall are the viral isolations (in the require reduced to the regular reduced to the regular reduced to the regular reduced to the regular reduced to the reduced to the reduced to the Artificial solution in December 1989 for Archieles and identificant Artificial response to the reduced to the reduced to the Artificial reduced to the reduced to the reduced to the reduced to the Artificial Reduced to the reduced t

But we use the expected speciment simples from the form weight we take the strong colors in resident out in example by the other than the converse of mean action and the strong of the  $C_{\bullet}$ 

TABLE I

# Virus Isolated in 1967 by Source and by Field Station

| Birds   |   |
|---|---|
| An 5417 - Myiobius atricaudus An 5560 - Conopophaga lineata An 5632 - Kanthomyas virescens An 5742 - Schiffornis virescens An 6109 - Dendrocoleptes platyrostris An 6142 - Dysithamnus mentalis An 6157 - Haplospiza unicolor An 6212 - Sporophila caerulescens An 6236 - Sittasomus griseicapillus An 6241 - Basileuterus auricapillus An 6291 - Platyrinchus mystaceus An 6789 - Turdus albicollis An 6829 - Emberizoides herbicola | - Casa Grande - 28/ 7/66 - Casa Grande - 19/ 8/66 - Itapetininga - 26/ 8/66 - Casa Grande - 1/ 9/66 - Itapetininga - 24/11/66 - Casa Grande - 1/12/66 - Casa Grande - 1/12/66 - Itapetininga - 3/12/66 - Casa Grande - 16/11/66 - Casa Grande - 16/12/66 - Itapetininga - 23/12/66 - Itapetininga - 23/12/66 - Casa Grande - 3/ 3/67 - Casa Grande - 10/ 3/67 |
| An 7126 - Platyrinchus mystaceus  | - Itapetining - 6/4/67  |
| An 4229 - Philander opossum  Mosquitoes *   | - Casa Grande - 14/1/66   |
| Ar 4253 - Psorophora ferox<br>Ar 6171 - Wyeomyia confusa<br>Ar 6629 - Gulex sp<br>Ar 6813 - A.(K.) cruzii   | - Cotia - 21/1/66<br>- Casa Grande - 2/12/66<br>- Casa Grande - 3/2/67<br>- Casa Grande - 9/3/67  |

<sup>\* =</sup> Pools

TABLE 11

Results obtained in HI and NT tests with Human Sera from 1966
, in Casa Grande \*

| VIRUS      | RESULTS | RESULTS VIRUS RES |                                 |
|------------|---------|-------------------|---------------------------------|
| Group ▲    | 0/242   | Bunyamwera Group  | 7/24                            |
| EEE        |         | Guaroa            |                                 |
| WEE        |         | Cache Valley      |                                 |
| Mayaro     |         | Ar 2984           | 7                               |
| Mucambo    |         | California        | 0/24                            |
| Group B    | 26/242  |                   |                                 |
| Bussuquara | 24      | Group Phlehotomus | 1/24                            |
| SLE        | 22      | Itaporanga        | ī                               |
| YF.        | 5       | Icoaracy          |                                 |
| Ilheus     | 1.5     | Tacaiuma          |                                 |
| Group C    | 8/242   | (SP Ar 2317)      | 47/24                           |
| Marituba   |         | Boraceia          |                                 |
| Caraparu   | 3       | SP Ar 395         | ns <b>(</b> n)                  |
| Oriboca    | 6       | Ar 43.75          | 25 <b>/</b> 24<br>9 <b>/</b> 24 |

<sup>\*</sup> Number of positives / Number of tested

TABLE III

Pools of Mosquitoes Inoculated in 1967

| SPECIES   | Cara<br>Grande                                      | Itapet.                                     | Cotia                                      | Total   |
|---|---|---|--|---|
| Anophelini Archeles cruzii k strodei utzi   | 108<br>4  | 1<br>1                                      | 1<br>18<br>3                               | 110<br>19<br>7  |
| Culicini  Aedes fluviatilis  leucocelaemus  serratus  crinifer  Culex (Melanoconion) sp  (Microculex) sp  Mansunia albifera  venezuelensis  titilans  Psorophora albipes  discrucians  ferox  Uranotaenia ditaenionota      | 1<br>3<br>14<br>2<br>2                              | 1<br>10<br>3<br>3<br>4<br>1<br>2<br>2<br>30 | 4<br>1<br>4<br>2<br>13<br>1<br>7<br>1<br>9 | 1 48 4 9 2 19 1 8 2 3 54 1                              |
| Sabethini Limatus flavisetosus Phoniomyla pilicauda Sabethes albiprivus intermedius Trichoprosopon digitatum pallidiventer reversum Wyeomyla confusa leucostigma Trichoposopron fluviatilis Others Dipterous Phlebotomus sp | 1<br>78<br>2<br>7<br>1<br>35<br>22<br>24<br>1<br>11 | 1<br>2<br>1<br>2<br>4                       | 4<br>3<br>1<br>1                           | 2<br>84<br>3<br>7<br>1<br>1<br>23<br>29<br>1<br>11<br>2 |
| Simulium auristriatum TOTAL   | 333   | 68  | 74   | 475   |

TABLE IV
Birds Netted From August 1966 to December 1967

| SPECIES   | Itapet.           | Casa<br>Grande          | Total                    |
|---|-------------------|-------------------------|--------------------------|
| Tinamidae<br>Crypturellus obsoletus   |                   | 1                       | 1                        |
| Accipitridae<br>Accipiter erythronemius   | 1                 |                         | ı                        |
| Falconidae<br>Micrastur ruficollis  |                   | 1                       | 1                        |
| Rallidae<br>Laterallus sp   | 1                 |                         | 1                        |
| Columbidae<br>Columbigallina talpacoti<br>Leptotila rufaxilla<br>Oreopeleia montana   | 69<br>7<br>4      | 6                       | 69<br>7<br>10            |
| Cuculidae<br>Dromococcym pavoninus<br>Crotophaga ani<br>Guira guira   | 4<br>5<br>2       |                         | 4<br>5<br>2              |
| <u>Psittacidae</u><br>Triclaria malachitacea  |                   | 1                       | 1                        |
| Strigidae<br>Otus choliba   | 2                 |                         | 2                        |
| Caprimulgidae<br>Hydropsalis brasilianus<br>Eleut. optus anomalis   | 1 1               |                         | 1                        |
| Trochilidae<br>Clytolasma rubricauda  |                   | 1                       | 1                        |
| Alcedinidae<br>Chloroceryle americana   | 7                 |                         | 7                        |
| Bucco ap  | 1                 |                         | 1                        |
| Picidae<br>Crysoptilus melanochlores<br>Picumus temminckii  | 1 6               |                         | 1 6                      |
| Dendrocolaptidae Dendrocolaptes platyrostris  squamatus Lepidoco uptes fuscus Campylorhamphus trochilirostris Sittasquus griseicapillus | 9<br>8<br>7<br>15 | 9<br>1<br>27<br>2<br>23 | 18<br>1<br>35<br>9<br>38 |

TABLE IV

Birds Netted from August 1966 to December 1967 - Cont.

| SPECIES                          | Itapet. | Casa<br>Grande                                 | Total                 |
|----------------------------------|---------|--|-----------------------|
| Filmariidae<br>Furnarius rufus   | 6       |  | 6                     |
| Synallaxis ruficapilla           | 10      | ,  | 13                    |
| Anabazenops fuscus               | 10      | 3<br>1<br>5<br>3<br>2<br>2<br>2<br>3<br>7<br>2 | 2                     |
| Syndactyla rufosuperciliata      | 25      | 5  | 30                    |
| Xenicopsoides amaurotis          | ~       | 1 3  |                       |
| Philyder atricapillus            | 2       | 1 3  | 3<br>2<br>2           |
| " rufus                          | ~       | 2  | 2                     |
| Automulus leucophthalmus         | 42      | 2  | 44                    |
| Cichlocolaptes leucophrys        | 4~      | 2  |                       |
| Heliobletus contaminatus         | 1       | 7  | Ŕ                     |
| Xenops minutus                   | 1 -     |  | 2                     |
| " rutilans                       | 1       | l ~ I  | 2                     |
| Sclerurus scansor                |         | 4  | 3<br>8<br>2<br>2<br>4 |
| Lochmias nematura                | 8       | 9  | 17                    |
|                                  |         | 1  | - '                   |
| <u>Formicariidae</u>             | Į.      | ]  |                       |
| Batara cinerea                   |         | 3  | 3<br>1                |
| Mackenziaena leackii             | 1       | 1  |                       |
| Thamnophilus caerulescens        | 22      | 8  | 30                    |
| " ruficapillus                   | 4       |  | 4                     |
| Dysitharms mentalis              | 13      | 16   | 29                    |
| Myrmotherula gularis             |         | 9  | 9                     |
| Drymophila malura                | 4 2     |  | 4                     |
| Pyriglena leucoptera             | 2       | 17   | 19                    |
| Chamaeza camparisoma             | }       | 1  | 1                     |
| Myracderus squamosus             |         | 5  | 5                     |
| Conopophagidae                   |         | •  |                       |
| Conopophagicaineata              | 55      | 20   | 75                    |
| <del>.</del>                     | 1       | 1  |                       |
| Rhinocryptidae<br>Merulaxis ater |         |  | ,                     |
| meritaxis ater                   | 1       | }  | 1                     |
| Cotingidae                       | ľ       | !  |                       |
| Attila rufus                     | 1       | 2  | 3                     |
| * phoenicurus                    | İ       | 6  | 6                     |
| Pachyramphus polychopterus       | 2       | 1  | 36332                 |
| Platypsaris rufus                | 3       | <b>f</b>                                       | 3                     |
| Frocnias nudicollis              | 1       | 1  | 2                     |
| Pipridae                         |         |  |                       |
| Piprites chloris                 | }       | ,  | ,                     |
| Chiroxiphia caudata              | 117     | 100  | 217                   |
| Ilicura militaris                | 111     |  | 217                   |
| Manacus manacus                  |         | 14   | 14                    |
| Schiffornis virescens            | סר      | 2  | 2                     |
| Neopelma aurifrons               | 18      | 28   | 46                    |
| neoberny gritiitong              | 2       | 26   | 28                    |

ρ

TAME IV

Birds Nettel From August 1966 to December 1967 - Cont.

| SPECIES   | Itapet.  | Casa<br>Graide                                   | Total   |
|---|--|--|---|
| Tyron Mass Satrapa icterophrys Muscivere tyrannus Tyrannus melancholicus Empidenomus varius Sirystes sibilatrix Pitengus sulphuratus Myiarchus tyrannulus Empidenax euleri Myiebius atricaudus Myiephebus fasciatus Platyrinchus mystaceus Telmenyias sulphurescens Tedrenyias sulphurescens Todirestrum policcephalum  m plumbeiceps Henetricus diops Phylloscurtes ventralis Serpophaga subcristata Elaenia chariquensis mesolence cristata Camptestoma otsoletum Leptopogon amaurecephalus Pigrenerpia rufiventris Himudirlane | 4<br>3<br>5<br>1<br>5<br>2<br>32<br>14<br>42<br>2<br>1<br>8<br>5<br>30<br>49<br>2<br>2 | 1<br>4<br>12<br>36<br>3<br>1<br>2<br>1<br>3<br>5 | 4 3 5 1 1 5 2 36 12 4 78 5 1 1 2 9 5 33 54 2 2 9 79 |
| Niconfices Puroprogram tapera fusca Stelfic ptergy rufficollis Alppochelidon fucata   | 1<br>11<br>1   |  | 1<br>11<br>1  |
| <u>Trontogytione</u><br>Tropicdytes musullus  | 6  |  | 6   |
| <u>Mindelae</u><br>Mirus asturninua   | 4  |  | 4   |
| <u>Pleceione</u><br>Payver concatieus   | 1.   |  | 1   |
| Turdings Turdus allicoilis " ansure helicus " louceseess " rufiventris Fatycichla flavipes  | 30<br>39<br>2<br>59  | 31.<br>3<br>2<br>24<br>32                        | 61<br>42<br>4<br>83<br>32                           |
| Cyclarhidae<br>Cyclarhidaeochre ng lawa   | 5  | 3  | 8   |

TABLE IV

Birds Netted From August 1966 to December 1967 - Cont.

| SPECIES                     | Itapet.           | Casa<br>Grande                   | Total               |
|-----------------------------|-------------------|----------------------------------|---------------------|
| Vireonidae                  |                   |                                  |                     |
| Vireo chivi                 | 18                | 2                                | 20                  |
| Hylophilus poicilotis       | 8                 | 8                                | 16                  |
| Coerebidae                  | 1                 |                                  | İ                   |
| Dacnis cayana               | 1 2               |                                  | 2 3                 |
| Coereba flaveola            | 2 3               | }                                | 3                   |
| Compacthlypidse             |                   |                                  |                     |
| Geothlypis aequinoctialis   | 3                 | <b>j</b>                         | 3                   |
| Besileuterus leucoblepharus | 3 8               | 5                                | 3<br>13<br>14<br>11 |
| hypoleucus                  | 14                |                                  | 14                  |
| " auricapillus              | 14                | 10                               | 11                  |
| Thraupidae                  | 1                 | [                                |                     |
| Tanagra pectoralis          | 1                 | 6                                | 6                   |
| Pipraeidea melanonota       | { <b>1</b>        | 6                                |                     |
| Tangara seledon             |                   | 4                                | 7<br>4<br>4<br>9    |
| " cyanocephala              | ľ                 | 4 9                              | 4                   |
| " desmaresti                |                   | 9                                | 9                   |
| " cayana                    | 14                | j i                              | 14                  |
| Thraupis cyanoptera         | 1                 | 13                               | 13                  |
| " sayaca                    | 32                | 1                                | 33                  |
| Orthogonys chloricterus     |                   | 13<br>1<br>3<br>5 <b>5</b><br>16 | 3<br>16             |
| Habia rubica                | 11                | 5 <b>5</b>                       | 16                  |
| Tachyphonus coronatus       | 65                |                                  | 81                  |
| Trichothraupis melanops     | 12                | 48                               | 60                  |
| Neothranpis fasciata        | l                 |                                  | 1                   |
| Schistochlamys melanopis    | 20                | 1                                | 20                  |
| <u>Icteridae</u>            | į                 | ] .                              |                     |
| Molothrus bonariensis       | 26                | Į                                | 26                  |
| Omorimopsar chopi           | 2                 |                                  | 2                   |
| Fringillidae                | ļ                 | 1                                |                     |
| Saltator similis            | 10                | 10                               | 20                  |
| Tiaris fuliginosa           | 1                 | ] ]                              | 2                   |
| Sporophila caerulescens     | 38                | 1                                | 38                  |
| * plumbea                   | 38<br>1<br>9<br>5 | (                                | 38<br>1<br>9<br>5   |
| Volatinia jacarina          | 9                 |                                  | ] 9                 |
| Spinus magellanicus         | 1 3               | ł                                |                     |
| Sicalis flaveola            | 6                 | 1 70                             | 6                   |
| Haplospiza unicolor         | 2                 | 19                               | 21                  |
| Arremon taciturnus          | 5                 | 1 +                              | 6                   |
| Myospiza humeralis          | 12                | (                                | 12                  |
| Zonotrichia capensis        | 149               | 1                                | 149                 |
| Emberizoides herbicola      | 7 2               | <b>\</b>                         | 7 2                 |
| Donacospiza albifrons       |                   | L                                |                     |
| TOTAL                       | 1 362             | 784                              | 2 146               |

TABLE \_\_V
Trapped
Mamanis Notice From August 1966 to December 1967

| SPECIES   |   | ASA<br>ANDE                         | ITAPET.                  | TOTAL                                    |
|---|---|-------------------------------------|--------------------------|--|
| Rodente   |   |                                     |                          |  |
| Muridae Rattus norvegicus rattus Mus musculus   |   | 35<br>22<br>1                       | 1<br>23                  | 36<br>45<br>1                            |
| Cricetidae Oryzemys nigripes  ** laticeps  ** subflavus  ** capito Delemys dersalis Nectomys squamipes Akoden srvieuleides Thaptomys nigrita Oxymyeterus quaester |   | 12<br>12<br>12<br>3<br>4<br>35<br>4 | 1<br>1.<br>19<br>19<br>1 | 12<br>2<br>1<br>12<br>3<br>23<br>54<br>5 |
| Erethizontidae<br>Coerdon insidiosus  |   | 1                                   |                          | 1  |
| Cavildae<br>Cavia aperea  |   | 3                                   |                          | 3  |
| Echinyidae<br>Proechinys iheringi<br>Chyonys laticeps<br>Not Identified   |   | 19<br>151                           | 10<br>20                 | 19<br>10<br>171                          |
| <u>Marsupials</u>   |   |                                     |                          |  |
| Didelphis marsuptallis Monodelphis tricolor Philander opossum   |   | なるた                                 | 3                        | 17<br>2<br>6                             |
| Not Inentified  | - |                                     |                          | ì  |
| Bats  |   |                                     |                          |  |
| Carrolia perspicilata Desmodus rotundus   |   | 2<br>1                              |                          | 2  |
| Not Identified  |   | n/                                  | 10                       | 104                                      |
| TOTAL   |   | 435                                 | 109                      | 544                                      |

Jnclassi ieu

| Security Classification  |  |   |   |
|--|--|---|---|
|  | NTROL DATA - RE  |   | the averall toward in algorithm?  |
| (Security classification of title, body of abstract and index  1. ORIGINATING ACTIVITY (Corporate author)  | ing annotation must be e   |   | RT SECURITY CLASSIFICATION  |
| Instituto Adolfo Eutz, Laboratório d   | a Amianiana  |   |   |
| São Paulo, Brasil  | e Arbovitus,   | 2 b GROUI   | Unclassified  |
| 318311   |  |   | · NA  |
| 3 REPORT TITLE   |  | <u> </u>  | <u> </u>  |
| ANGOVERNO STUDIES EN SÃO TANDA, ER   | MIZL.  |   | •   |
| 4. DESCRIPTIVE NOTES (Type of report and inclusive dates)  |  |   |   |
| Annual Technical eport, period llo   | vember 1966  | January   | 1967  |
| Lopes, Oscar de Souza  |  |   |   |
| 6 REPORT DATE  | 78 TOTAL NO. OF  | PAGES   | 76. NO. OF REFS   |
| January 1967   | 12   |   |   |
| BE CONTRACT OR GRANT NO.   | 98. ORIGINATOR'S R   | EPORT NUM   | BER(S)  |
| DfHC19-67-7-0010<br>b. project no.   |  |   |   |
| 2H0145C1371D 00 020 17.  | Ì  |   |   |
| e.   | 9b. OTHER REPORT   | NO(S) (Any  | other numbers that may be assigned  |
| d.   |  |   |   |
| 10 AVAILABILITY/LIMITATION NOTICES   | <del>*************************************</del>   |   |   |
| Unlimited  |  | <u></u>   |   |
| 11. SUPPLEMENTARY NOTES  | 12 SPONSORING MIL<br>U.S. Azmy 31  |   | vity<br>lefense Research Cifice   |
| None   | Latin Amer   | ica   | ा York 09676  |
| 13. ABSTRACT Reports on continuation of birds, animals, and human sera have to detected. A few new, or new isolation yale Arbovirus Laboratory for final it isolations were also made from pooled of houses, and inside of houses. Conspeciments from ill inhabitants retice to indicate transmission from lorest concept, evidence, remaining problems. | peen studied is one in the studied in the studied identification is mosquitos colutional difficular to report in reservoirs in the second in t | olating<br>y area h<br>or chara<br>lected i<br>lty, expe<br>liness.<br>o houses | and screening viruses have been sent to the acterization. In the forest, outside crienced in obtaining Swidence continues by wild mosquitoes. |
| •• , ,   |  |   |   |

DD 5984. 1473

Unclassified
Security Classification

### Unclassified Security Classification

| KEY WORDS                               | LIP                                   | NK A | LINK B |      | LIN | кc   |      |
|---|---------------------------------------|------|--------|------|-----|------|------|
|   | · · · · · · · · · · · · · · · · · · · | ROLE | wt     | ROLE | wt  | ROLE | . ΑΤ |
| Virus                                   | •                                     |      |        |      |     |      |      |
| H <b>emorrha</b> gi.<br>Bpidemiolo      | gу                                    |      |        |      | 1   |      |      |
| ∴achupo Vii<br>Junin /irus<br>Arbovirus | rus<br>s                              |      | !<br>! |      |     |      |      |
|   | ·                                     |      |        |      |     |      |      |
|   |                                       |      |        |      |     |      |      |
|   |                                       | }    |        |      | •   |      |      |

- ORIGINATING ACTIVITY: Enter the name and address of the contractor, subcontractor, grantee, Department of Defense activity or other organization (corporate author) issuing the report.
- 2a. REPORT SECURITY CLASSIFICATION: Enter the overall security classification of the report. Indicate whether "Restricted Data" is included. Marking is to be in accordance with appropriate security regulations.
- 2b. GROUP: Automatic downgrading is specified in DoD Directive 5200.10 and Armed Forces Industrial Manual. Enter the group number. Also, when applicable, show that optional markings have been used for Group 3 and Group 4 as authorized.
- 3. REPORT TITLE: Enter the complete report title in all capital letters. Titles in all cases should be unclassified. If a meaningful title cannot be selected without classification, show title classification in all capitals in parenthesis immediately following the title.
- 4. DESCRIPTIVE NOTES: If appropriate, enter the type of report, e.g., interim, progress, summary, annual, or final. Give the inclusive dates when a specific reporting period is covered.
- 5. AUTHOR(S): Enter the name(s) of author(s) as shown on or in the report. Enter last name, first name, middle initial. If military, show rank and branch of service. The name of the principal author is an absolute minimum requirement.
- REPORT DATE: Enter the date of the report as day, month, year, or month, year. If more than one date appears on the report, use date of publication.
- 7a. TOTAL NUMBER OF PAGES: The total page count should follow normal pagination procedures, i.e., enter the number of pages containing information.
- 76. NUMBER OF REFERENCES: Enter the total number of references cited in the report.
- 8a. CONTRACT OR GRANT NUMBER: If appropriate, enter the applicable number of the contract or grant under which the report was written.
- 86, 8c, & 8d. PROJECT NUMBER: Enter the appropriate military department identification, such as project number, subproject number, system numbers, task number, etc.
- 9a. ORIGINATOR'S REPORT NUMBER(S): Enter the official report number by which the document will be identified and controlled by the originating activity. This number must be unique to this report.
- 9b. OTHER REPORT NUMBER(S): If the report has been assigned any other report numbers (either by the originator or by the sponsor), also enter this number(s).

- 10. AVAILABILITY LIMITATION NOTICES: Enter any limitations on further dissemination of the report, other than those imposed by security classification, using standard statements such as:
  - "Qualified requesters may obtain copies of this report from DDC."
  - (2) "Foreign announcement and dissemination of this report by DDC is not authorized."
  - (3) "U. S. Government agencies may obtain copies of this report directly from DDC. Other qualified DDC users shall request through
  - (4) "U. S. mulitary agencies may obtain copies of this report directly from DDC. Other qualified users shall request through
  - (5) "All distribution of this report is controlled. Qualified DDC users shall request through

If the report has been furnished to the Office of Technical Services, Department of Commerce, for sale to the public, indicate this fact and enter the price, if known

- 11. SUPPLEMENTARY NOTES: Use for additional explanatory notes.
- 12. SPONSORING MILITARY ACTIVITY: Enter the name of the departmental project office or laboratory sponsoring (paying for) the research and development. Include address.
- 13. ABSTRACT: Enter an abstract giving a brief and factual summary of the document indicative of the report, even though it may also appear e-sewhere in the body of the technical report. If additional space is required, a continuation sheet shall be attached.

It is highly desirable that the abstract of classified reports be unclassified. Each paragraph of the abstract shall end with an indication of the military security classification of the information in the paragraph, represented as (TS), (S), (C), or (U).

There is no limitation on the length of the abstract. However, the suggested length is from 150 to 225 words.

14. KEY WORDS: Key words are technically meaningful terms or short phrases that characterize a report and may be used as index entries for cataloging the report. Key words must be selected so that no security classification is required. Idenfiers, such as equipment model designation, trade name, military project code name, geographic location, may be used as key words but will be followed by an indication of technical context. The assignment of links, rules, and weights is optional.

Unclassified